

Title	The invertible Toeplitz operators on the Bergman spaces (General topics on applications of reproducing kernels)
Author(s)	米田, 力生
Citation	数理解析研究所講究録 (2016), 1980: 152-158
Issue Date	2016-01
URL	http://hdl.handle.net/2433/224443
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

The invertible Toeplitz operators on the Bergman spaces

小樽商科大学 (Otaru University)

米田 力生 (Rikio Yoneda)

Abstract

In this paper, we study the invertible (and Fredholm) Toeplitz operators T_φ on the Bergman spaces with harmonic symbol.

Key Words and Phrases : Bergman spaces, Toeplitz operator, closed range, invertible operator, Fredholm operator.

Let D be the open unit disk in complex plane C . For $z, w \in D$, and $0 < r < 1$, let $\varphi_w(z) = \frac{w-z}{1-\bar{w}z}$ and $\rho(z, w) = \left| \frac{w-z}{1-\bar{w}z} \right|$ and $D(w, r) = \{z \in D, \rho(w, z) < r\}$.

Let $H(D)$ be the space of all analytic functions on D .

The space $L^2(dA(z))$ is defined to be the space of Lebesgue measurable functions f on D such that

$$\|f\|_{L^2(dA(z))} = \left\{ \int_D |f(z)|^2 dA(z) \right\}^{\frac{1}{2}} < +\infty,$$

where $dA(z)$ denote the area measure on D . The Bergman space $L_a^2(dA(z))$ is defined by

$$L_a^2(dA(z)) = H(D) \cap L^2(dA(z)).$$

For $\varphi \in L^2(dA(z))$, the Toeplitz operator T_φ with symbol φ is defined on $L_a^2(dA(z))$ by

$$T_\varphi f = P(\varphi f),$$

where $P(f)(z) = \int_D \frac{f(w)}{(1 - \bar{w}z)^2} dA(w)$.

Let X, Y be Banach spaces and let T be a linear operator from X into Y . Then T is called to be bounded below from X to Y if there exists a positive constant $C > 0$ such that $\|Tf\|_Y \geq C \|f\|_X$ for all $f \in X$, where $\|\cdot\|_X, \|\cdot\|_Y$ be the norm of X, Y , respectively. The Berezin transform of T_φ is given by $\tilde{\varphi}(z) = \widehat{T_\varphi}(z) = \langle T_\varphi k_z, k_z \rangle$, where $k_z(w) = \frac{1 - |z|^2}{(1 - z\bar{w})^2}$.

If H is a Hilbert space, then a bounded operator T is a Fredholm operator if and only if the range of T is closed, $\dim \ker T$, and $\dim \ker T^*$ is finite.

For $a, b \in \mathbb{C}, \varphi, \psi \in L^\infty(D)$, then

- (a) $T_{a\varphi+b\psi} = aT_\varphi + bT_\psi$,
- (b) $T_{\bar{\varphi}} = T_\varphi^*$,
- (c) $T_\varphi \geq 0$ ($\varphi \geq 0$).

For $\varphi \in H^\infty$, then

- (d) $T_\psi T_\varphi = T_{\psi\varphi}$,
- (e) $T_{\bar{\varphi}} T_\psi = T_{\bar{\varphi}\psi}$

Let $\tilde{\varphi}$ denote the harmonic extension of the function φ to the open unit disk D . In [8], Douglas posed the following problem :

If φ is a function in L^∞ for which $|\varphi| \geq \delta > 0, z \in D$, then is T_φ invertible ?

And V.A. Tolokonnikov gave the following:

If $|\tilde{\varphi}(z)| \geq \delta > \frac{45}{46}$, then T_φ is invertible.

In [18], N.K.Nikolskii gave the following:

If $|\tilde{\varphi}(z)| \geq \delta > \frac{23}{24}$, then T_φ is invertible.

In [20], T.H.Wolff gave the following:

If $\inf_D |\tilde{\varphi}(z)| > 0$ and then T_φ is not invertible.

The study of Toeplitz operators on the Bergman spaces and Hardy space have been studied by many authors. In this paper, we study when the Toeplitz operators T_φ on the Bergman spaces with harmonic symbol is invertible or Fredholm.

In [14], the following theorem are well-known.

Theorem A. *Suppose φ is a bounded and nonnegative function. Then the following conditions are equivalent :*

- (1) T_φ is bounded below.
- (2) There is a constant $C > 0$ such that

$$\int_D |f(z)|^2 \varphi(z) dA(z) \geq C \int_D |f(z)|^2 dA(z),$$

for all $f \in L_a^2(dA(z))$.

In [14], D.Leucking proved the following results.

Theorem B. *Let $\alpha > -1$ and $p > 0$. Then the following are equivalent:*

- (1) There is a constant $C > 0$ such that

$$\int_D |f(z)|^p dA(z) \leq C \int_G |f(z)|^p dA(z)$$

for all $f \in L_a^p(dA(z))$

- (2) There is a constant $C > 0$ such that

$$\int_D |f(z)|^p (1 - |z|^2)^\alpha dA(z) \leq C \int_G |f(z)|^p (1 - |z|^2)^\alpha dA(z)$$

for all $f \in L_a^p((1 - |z|^2)^\alpha dA(z))$

- (3) For any $a \in D$ a subset G of D satisfy the condition that there exist $\delta > 0$ and $r > 0$ such that $\delta |D(a, r)| \leq |D(a, r) \cap G|$, where $|D(a, r)|$ is the (normalized) area of $D(a, r)$.

Theorem C. *Let φ be a bounded measurable function on D . Then there is a constant $\epsilon > 0$ such that*

$$\int_D |\varphi(z) f(z)|^p dA(z) \geq \epsilon \int_D |f(z)|^p dA(z)$$

for all $f \in L_a^p(dA(z))$ if and only if there exists $r > 0$ such that the set $\{z \in D : |\varphi(z)| > r\}$ satisfies condition (3) of Theorem 3.

Theorem D. *Let φ be a bounded positive measurable function on D . Then T_φ is invertible if and only if there exists $r > 0$ such that the set $\{z \in D : |\varphi(z)| > r\}$ satisfies condition (3) of Theorem 3.*

The following theorem is well-known(see [21]).

Theorem E. *Suppose that $\varphi \in \mathcal{C}(\overline{D})$. Then the following conditions are equivalent :*

- (1) T_φ is Fredholm.
- (2) φ is nonvanishing on the unit circle.

Theorem 1. *Let $g \in H^\infty$. Then the following are equivalent:*

- (1) $T_{\widetilde{g}}$ is invertible operator on $L_a^2(dA(z))$
- (2) T_g is invertible operator on $L_a^2(dA(z))$
- (3) $\inf_{z \in D} |\widetilde{T_{\widetilde{g}}}(z)| = \inf_{z \in D} |g(z)| > 0$

The problem which we must consider next is following.

Problem. *Let $g, h \in H^\infty$ and $g, h \in \mathcal{C}(\overline{D})$. Then the following are equivalent:*

- (1) $T_{g+\overline{h}}$ is invertible operator on $L_a^2(dA(z))$
- (2) $\inf_{z \in D} |\widetilde{T_{g+\overline{h}}}(z)| = \inf_{z \in D} |g(z) + \overline{h(z)}| > 0$

At first, we can prove the following.

Theorem 2. *Let $g \in H^\infty$ and $g \in \mathcal{C}(\overline{D})$. Then the following are equivalent:*

- (1) $T_{g+\overline{g}}$ is bounded below on $L_a^2(dA(z))$
- (2) $T_{g+\overline{g}}$ is invertible operator on $L_a^2(dA(z))$
- (3) $\inf_{z \in D} |\widetilde{T_{g+\overline{g}}}(z)| = \inf_{z \in D} |g(z) + \overline{g(z)}| > 0$

Next, we can prove the following.

Theorem 3. *Let $g \in H^\infty$ and a constant $c > 1$. Then the following are equivalent:*

- (1) T_g is bounded below on $L_a^2(dA(z))$
- (2) $T_{cg+\bar{g}}$ is bounded below on $L_a^2(dA(z))$

Using Theorem 3, we prove the following main result.

Theorem 4. *Let $g \in H^\infty$ and $g \in \mathcal{C}(\overline{D})$. Then the following are equivalent:*

- (1) $T_{\bar{g}}$ is invertible operator on $L_a^2(dA(z))$
- (2) T_g is invertible operator on $L_a^2(dA(z))$
- (3) $T_{cg+\bar{g}}$ is invertible operator on $L_a^2(dA(z))$ ($c > 0, c \neq 1$)
- (4) $\inf_{z \in D} |\widetilde{T_{cg+\bar{g}}}(z)| = \inf_{z \in D} |cg(z) + \overline{g(z)}| > 0$ ($c > 0, c \neq 1$)
- (5) $\inf_{z \in D} |\widetilde{T_g}(z)| = \inf_{z \in D} |\widetilde{T_{\bar{g}}}(z)| = \inf_{z \in D} |g(z)| > 0$

Moreover, we can prove the following.

Theorem 5. *Let $g, h \in H^\infty$ with $\inf_{z \in D} |h(z)| - \sup_{z \in D} |g(z)| > 0$.*

If $\inf_{z \in D} |\widetilde{T_{g+\bar{h}}}(z)| > 0$, then $T_{g+\bar{h}}$ is invertible on $L_a^p(dA(z))$, and $T_{h+\bar{g}}$ is invertible on $L_a^p(dA(z))$

Theorem 6. *Suppose that $g \in H^\infty$ and $g \in \mathcal{C}(\overline{D})$. Then the following conditions are equivalent :*

- (1) $T_{\bar{g}}$ is Fredholm.
- (2) $T_{cg+\bar{g}}$ is Fredholm ($c > 0, c \neq 1$).
- (3) g is nonvanishing on the unit circle.

References

- [1] A.Aleman and A.G.Siskakis, Integration operators on Bergman spaces, Indiana Univ.Math.J.46(1997),337-356.
- [2] S.Axler, The Bergman space, the Bloch space, and commutators of multiplication operators, Duke Math.J.53(1986),315-331.
- [3] C.J.Bishop, Bounded functions in the little Bloch space, Pacific J.Math.142(1990), no.2,209-225.

- [4] Paul S.Bourdon, Similarity of parts to the whole for certain multiplication operators, *Proc.Amer.Math.Soc.*99(1987),563-567.
- [5] Paul S.Bourdon and J.H.Shapiro, Spectral Synthesis and Common Cyclic vectors, *Michigan Math.J.*37(1990), 71-90.
- [6] J.Bonet, P.Domanski, M.Lindstrom, Pointwise multiplication operators on weighted Banach spaces of analytic function, *Studia Math.* 137(1999),177-194.
- [7] L.Carleson, An interpolation problem for bounded analytic functions, *Amr.J.Math.* 80(1958),921-930.
- [8] R.G.Douglas, *Banach Algebra Techniques in Operator Theory*, Academic Press, New York, 1972.
- [9] P.Duren, A.Schuster , *Bergman Spaces*, American Mathematical Society, Providence, 2004.
- [10] P.Ghatage and D.Zheng and Nina Zorboska, Sampling sets and closed range composition operators on the Bloch space, *Proceedings of the Amer.Math.Soc.*133, 5(2004), 1371-1377.
- [11] J.B.Garnett and D.Marshall, *Harmonic Measure*, Cambridge University Press, 2005.
- [12] J.Bonet, P.Domanski, M.Lindstrom, Pointwise multiplication operators on weighted Banach spaces of analytic function, *Studia Math.* 137(1999),177-194.
- [13] H.Hedenmalm and B.Korenblum and K.Zhu, *Theory of Bergman Spaces*, Springer-Verlag, New York.
- [14] P.Jones, Ratios of Interpolating Blaschke products, *Pacific J.Math.*95(2) (1981), 311-321.
- [15] D.Leucking, Inequalities on Bergman spaces, *Illinois J.Math.*25(1981), 1-11.
- [16] D.Marshall, Blaschke products generate H^∞ , *Bull Amer.Math.Soc.*, 82(1976), 494-496.
- [17] G.McDonald and C.Sundberg, Toeplitz operators on the disc, *indiana Univ.Math.J.* 28(1979),595-611.
- [18] N.H.Nikolskii, *Treatise of the Shift Operator*, Springer-Verlag, Berlin, 1986.
- [19] K.Seip, Beuring type density theorems in the unit disk, *Invent.Math.*113(1993), 21-39.
- [20] T.H.Wolff, Counterexamples to two variants of the Helson-Szego theorem, Report No.11, California Institute of Technology, Pasadena, 1983.

- [21] R.Zhao, On α -Bloch functions and VMOA, Acta Math.Sci.16(1996), 349-360.
- [22] K.Zhu, Multipliers of BMO in the Bergman metric with applications to Toeplitz operators, J.Funct.Anal. 83(1989),31-50.
- [23] K.Zhu, Operator Theory in Function Spaces, American Mathematical Society, Providence, 2007.
- [24] K.Zhu, Bloch type spaces of analytic functions, Rocky Mout.J.Math. 23(1993), 1143-1177.